

Efficacy Information

SKU.1020 - Clinical Midi Wipes 200 Count – Formulation Efficacy

The efficacy as shown below has been independently tested.

Study Title	Target Organism	Result
EN 13727:2015 Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of bactericidal activity in the medical area - Test method and requirements (phase 2, step 1)	Staphylococcus aureus Enterococcus hirae Pseudomonas aeruginosa	log reduction: ≥ 5 log ₁₀ Pass under dirty conditions
EN 1276 Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of bactericidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic and institutional areas – Test method and requirements, (phase 2, step 1)	Pseudomonas aeruginosa Escherichia coli Staphylococcus aureus Enterococcus hirae	log reduction: ≥ 5 log ₁₀ ; Pass under dirty conditions
EN 14476:2005 Chemical disinfectants and antiseptics - Virucidal quantitative suspension test for chemical disinfectants and antiseptics used in human medicine. Test method and requirements (phase 2, step 1)	Influenza A (H1N1) (TC Adapted) (ATCC- VR-1469)/MDCK cells (ATCC-CCL-34)	log reduction: ≥ 3 log ₁₀ ; Pass under dirty conditions
	Hepatitis C Virus (Bovine Viral Diarrhoea Virus surrogate) ATCC VR – 1422 BT Cells	log reduction: ≥ 4 log ₁₀ ; Pass under dirty conditions
	Herpes Simplex Virus 1 (ATCC VR-733)/VERO cells (ATCC CCL-81)	log reduction: ≥ 4 log ₁₀ ; 20 °C Pass under dirty conditions
	Human norovirus surrogate (Feline calicivirus/CRFK cells)	log reduction: ≥ 4 log ₁₀ ; Pass under dirty conditions